





Chemical Abundances

Equivalent Width




$\int I_{\lambda} d\lambda$ at two different wavelengths?

Same total energy? Same number of absorbing atoms?



Line profile $\phi_\nu = \frac{I_c - I_\lambda}{I_c}$



$$W_\lambda = \int_{-\infty}^{\infty} \frac{I_c - I_\lambda}{I_c} d\lambda = \int (1 - e^{-\tau_\nu}) d\lambda$$

measures the total absorption (strength) in a line, where I_c is the continuum and I_λ is the line profile.

W_λ has the dimension of λ , e.g., Å, or mÅ.

In optical and UV ($h\nu \gg kT$), stimulated emission can be neglected, i.e., $1 - \exp(-h\nu/kT) \rightarrow 1$, $\sigma_\nu \rightarrow \sigma_0$

Curve of Growth

... used in estimation of the abundance of a species by the equivalent width of a spectral line

$$\tau_\nu = \kappa_\nu ds = n\sigma_\nu ds = N \sigma_\nu$$

where N is the column density

$$\sigma_\nu = \left(\frac{\pi e^2}{mc}\right) f \phi_\nu, \quad \sigma_\nu d\nu = \sigma_\lambda d\lambda$$

$$\tau_\lambda = N \left(\frac{\pi e^2}{mc^2}\right) f \lambda_0^2 \phi_\lambda$$

- (i) For weak lines ($\tau_\lambda \ll 1$) Every absorbing atom/ion contributes to W

$$W_\lambda = \int \tau_\lambda d\lambda = N \frac{\pi e^2}{mc^2} f \lambda_0^2 \propto N f$$

or

$$\frac{W_\lambda}{\lambda} = N \frac{\pi e^2}{mc^2} f \lambda_0 = 8.85 \times 10^{-13} N f \lambda$$

where N is in $[\text{cm}^{-2}]$, and λ in $[\text{cm}]$.



(ii) For strong lines ($\tau_\lambda \gg 1$) **Line wings**

$$W_\lambda \propto \sqrt{N f}$$

(iii) Intermediate case

$$W_\lambda \propto \sqrt{\ln N f}$$

Line starts to “saturate”, W almost independent of number of absorbing atoms






Figure 9.22 A general curve of growth for the Sun. (Figure from Aller, *Atoms, Stars, and Nebulae*, Revised Edition, Harvard University Press, Cambridge, MA, 1971.)

- W_λ gives the abundance of the absorbing atom/ion.
- It is a coarse estimate ($\sim 5\%$), with many uncertainties. A more accurate measurement requires high-dispersion spectroscopy, stellar atmosphere modeling, etc.
- If it is an excitation transition, use the Boltzmann equation to compute the total number of the species.
- Then use the Saha equation to compute the total number of the element.

INTERSTELLAR ABSORPTION LINES IN THE SPECTRUM OF ZETA OPHIUCHI

DONALD C. MORTON

Princeton University Observatory

Received 1974 August 12

ABSTRACT

Extensive high-resolution scans with the ultraviolet spectrometer on the *Copernicus* satellite have been combined with the available ground-based data on interstellar lines to obtain temperatures, densities, and abundances in H I clouds and H II regions in the direction of ζ Oph. Column densities have been obtained for 21 elements in various stages of ionization. A new determination for CO and the results for H₂, HD, CH, CH⁺, and CN for other authors also have been included. In addition, upper limits have been reported for five elements and 11 molecules. In the ultraviolet scans, 45 lines remain unidentified. Radial velocities and curves of growth were used to locate the species seen in the ultraviolet into one or more of the six clouds already known from the visible spectra. The H₂, HD, and most of the neutral atoms are concentrated in one cloud at a heliocentric velocity of -14.4 km s^{-1} , while N I, O I, Ar I, and most of the first ions are distributed over at least two of these clouds. The velocities of the higher ion states implied there are H II regions in addition to the Strömgren sphere.

Calculations of the ionization equilibrium for C, Mg, S, and Ca have shown that the electron density $n_e \sim 0.7 \text{ cm}^{-3}$ in the -14.4 km s^{-1} cloud. Since the ionization of carbon is the primary source of the electrons, the ratio C II/H requires that $n_H \sim 10^4 \text{ cm}^{-3}$ for the total hydrogen nuclei, with a fractional ionization $n_e/n_H \sim 7 \times 10^{-5}$. Both densities could be up to 20 times larger if the cloud is close to the star. The cloud must be no more than 0.05 pc thick. The populations of the fine-structure levels in the C I ground state require $T = 19^\circ \text{ K}$. The HD and the excited rotational levels of H₂ need temperatures between 56° and 115° , though the lines appear to have the same velocity as the C I, probably indicating that the dense cloud must have some associated hotter regions.


Relative to hydrogen, most of the elements in the H I clouds are depleted by factors of 3 to 4000 compared with the solar system abundances. Only sulfur and zinc are present in the gas with near normal abundances. Several elements, particularly Al, Si, and Fe, appear to be depleted in the H II regions as well, but nitrogen is normal.

Zeta Ophiuchi = HD 149757, O9.5V (*so few stellar lines*), a rapid rotator $v \sin i \approx 250$ km/s (*so broad stellar lines* \rightarrow *easy separation from the narrow interstellar lines*) by Cope

Ex 1:

K I line $\lambda 7699$, $f = 0.339$, W_λ measured = 84 mÅ $\rightarrow N(\text{K I}) = ?$

Ex 2: Application to unidentified lines the “diffuse interstellar bands” (DIBs), 100+ such bands most between 440 to 685 nm. Ref: Herbig 1995, ARAA, 33, 359




Do these lines originate from abundances?

$$I = I_0 e^{-\tau} = I_0 e^{-N\sigma}$$

Observed $I/I_0 = 0.99$

$\Delta\lambda \approx 25$ Å, extremely broad for Doppler broadening

\rightarrow estimate N_x , by assuming extreme values of f and compare with N_H



Ex 3: The Doublet Ratio Method \rightarrow 2 lines from the same elements, e.g., close doublets Na I, Ca II \leftarrow Saha eq.

Abundances of Elements

In HI regions, $N(x)/N(H)$ more or less depleted with respect to atmospheres of Pop I stars

(1) into molecular forms? If so, depletion should occur primarily in regions of high densities

Results: Indeed, depletion seen where $E_{B-V} > 0.3$

For $E_{B-V} < 0.05$, abundances normal

(2) into solid forms? If so, depletion should increase for elements of higher condensation temperatures

Results: Indeed, this was observed

$T_c \uparrow \rightarrow$ condense first \rightarrow more depletion

INTERSTELLAR ABUNDANCES: GAS AND DUST

GEORGE B. FIELD

Center for Astrophysics, Harvard College Observatory and Smithsonian Astrophysical Observatory

Received 1973 August 8

ABSTRACT

Data on abundances of interstellar atoms, ions, and molecules in front of ζ Oph are assembled and analyzed. The gas-phase abundances of at least 11 heavy elements are significantly lower, relative to hydrogen, than in the solar system. The abundance deficiencies of certain elements correlate with the temperatures derived theoretically for particle condensation in stellar atmospheres or nebulae, suggesting that these elements have condensed into dust grains near stars. There is evidence that other elements have accreted onto such grains after their arrival in interstellar space. The extinction spectrum of ζ Oph can be explained qualitatively and, to a degree, quantitatively by dust grains composed of silicates, graphite, silicon carbide, and iron, with mantles composed of complex molecules of H, C, N, and O. This composition is consistent with the observed gas-phase deficiencies.




FIG. 1.—Logarithmic abundances of gas-phase elements in ζ Oph, relative to solar-system values, plotted against the condensation temperature calculated for an oxygen-rich atmosphere (table 3). It is suggested that elements above 500° K condensed as cores in stellar nebulae or atmospheres, while those below 500° K condensed as mantles in interstellar space. The large deficiencies of C and Ca are consistent with the latter process.

ULTRAVIOLET STUDIES OF THE INTERSTELLAR GAS

Lyman Spitzer, Jr. and Edward B. Jenkins

Princeton University Observatory, Princeton, New Jersey 08540

(1975) ARAA




Figure 1 High-resolution scan of the O9.5 V star ζ Oph [$m_V = 2.56$, $E(B-V) = 0.32$] over an 8-Å interval. Error bars show the dispersion in photon counts expected from statistical fluctuations. The wavelengths of an Ar I line and of various rotational features in the (4,0) and (5,0) vibrational Lyman bands of H₂ and HD are shown by vertical lines.




Figure 2 Curves of growth for different groups of interstellar lines in ζ Oph. The filled circles represent lines produced by N I, Ar I, Mg II, Si II, S II, and Fe II; the triangles show C I, Na I, Mg I, S I, K I, and Fe I. The crosses represent H_2 Lyman lines from the rotational levels $J = 3$ – 6 .




Figure 5 Depletion below solar abundances for elements in their atomic form in the H I gas toward ζ Oph, plotted against atomic mass A in (a) and condensation temperature T_c in (b). The vertical width of each bar represents the experimental error arising from uncertainties in the respective element's curve of growth. For grains and atoms of a given charge, the nonequilibrium accretion rate in cool interstellar clouds should be proportional to $A^{-1/2}$, illustrated by the dotted line in (a). All elements shown here except N, O, and Ar should be predominantly ionized in H I regions.

ABUNDANCES OF INTERSTELLAR ATOMS FROM ULTRAVIOLET ABSORPTION LINES

EDWARD B. JENKINS
Princeton University Observatory

BLAIR D. SAVAGE
Washburn Observatory, University of Wisconsin

AND

LYMAN SPITZER, JR.
Princeton University Observatory

THE ASTROPHYSICAL JOURNAL, 301:355-379, 1986 February 1







FIG. 5.—Abundance of Fe vs. average density. Logarithm of the abundance ratios, $N(\text{Fe II})/N_{\text{H}}$, is plotted against the value of $\langle n_{\text{H}} \rangle \equiv N_{\text{H}}/R$, the average particle density of hydrogen, $[n(\text{H I}) + 2n(\text{H}_2)]$, along the line of sight of length R . The diameter of each circle is inversely proportional to the expected 1σ observational error in the logarithmic abundance ratio, as shown by the error bars in the legend. Solid curve represents a two-parameter fit to the theoretical result expected if the abundance ratio has one value in warm neutral gas and another in cold clouds. Note that $\log [N(\text{Fe})/N_{\text{H}}]_{\text{cosmic}} = -4.5$.



Estimated proportions of matter, dark matter and dark energy in the universe. Only the fraction of the mass and energy in the universe labeled "atoms" is composed of chemical elements. (*WMAP*)

<http://map.gsfc.nasa.gov/media/080998/index.html>



- ❑ What is the 1st, 2nd, 3rd, and 4th most abundant elements?
- ❑ What happens to iron?