




- A star → **a point source**
→ flux/magnitude
e.g., $m_v=15.7$
- A galaxy or central part of a globular cluster
→ **an extended source**
→ integrated flux, or surface brightness
e.g., 18.2 mag/sq arcsec
- The sky is an extended source.
In a dark site,
sky ~ 20 -21 mag/sq arcsec






Detectors


Eye as a detector

Pupil diameter

- determines resolving and light-gathering power
- adapts to existing light levels
- 8 mm (age 20) to 2.5 mm (age 80)



視網膜 The Retina



Light Sensitive Retinal Cells

- **rods** 視網膜桿

night vision $\lambda_{\max} \sim 507 \text{ nm}$

$V_{\text{lim}} \sim 8 \text{ mag}$

in practice 5.5-6.5 mag



outer periphery of retina

→ averted vision

- **cones**


color vision $\lambda_{\max} \sim 555 \text{ nm}$

central retina=fovea




Photography

- **Emulsion (感光乳劑)** :
micron-sized grains of some
silver halide (鹵化銀), e.g.,
silver bromide (溴化銀 ;
AgBr), suspended in a thin
layer of gelatin
- In astronomical applications,
add a glass base for support →
photographic plate



Photographic Process

- Photon → grain
→ e^- excited
+ Ag^+ (e.g., from thermal excitation)
→ $Ag + Ag + Ag \dots$
→ **latent image**
+ reducing agent (adding hydrogen or
removing oxygen or, in this case,
removing bromine)
→ conversion of entire grain to pure silver
- **Stop** once a clear image is obtain




Developing (顯影) 加大分開效果

Stop (急制)

Fixing (定影) 除去 Br 及剩下的 AgBr

- There is always some partially-developed grains remain → a 'fog' on the film



‘Reciprocity failure’ --- increasing inefficiency of photographic emulsions with longer exposure times

Note Most photoelectrons do not survive long enough to meet with a silver ion
→ **process very inefficient**
e.g., ~1000 photons → 1 developed grain

Quantum Efficiency (QE)

= The efficiency a device records the incident photons
= [# of records] / [# of incoming photons]

Photographic plates QE ~ a few %,
at best < 10%

Note To increase the sensitivity, some measures can be taken

- Adding chemical sensitizers to the emulsion
- Baking or soaking in nitrogen or hydrogen


→ Process of **hyersensitization**

Note In addition to **low sensitivity**, another disadvantage of a photographic plate is **non-linearity**


→ a given exposure may be correct for some stars, but overexposed for bright stars and underexposed for fainter stars

Even for a correctly exposed star, the density of the image does not directly reflect the brightness of a star, because of internal scattering

Incident photons are scattered within emulsion before being absorbed \rightarrow enlarged, circular image with size \sim # of scattering



Brightness of a star: a complicated function of total density AND image size



Photography --- Summary

Disadvantages

- low sensitivity
- complex procedures, thus prone to error
- density --- rather than intensity --- recorded

Advantages

- cheapness
- long exposure (cf. eye $\sim 0.1\text{s}$)
- ease of storage
- familiar techniques
- large field of view (suitable for observations)

Field of View

- Schmidt telescopes use photographic plates
30 cm square, corresponding to a sky area
 $\sim 6^\circ \times 6^\circ$
- Some plates are 50 cm square
- In comparison, modern electronic detectors
 $< 5\text{ cm}$ (1° FOV) \rightarrow mosaic


Plate Scale

$$P \text{ ["/pixel]} = (206,265) \times \overset{\text{CCD}}{\underset{|}{\mu}} \text{[micron]} / \overset{\text{Primary}}{\underset{|}{1,000}} \times f \text{ [mm]}$$

LOT (Lulin One-meter Telescope)

- $D=1000$ mm
- $f/8$
- E.g., CCD camera has
24 micron x 1024 x 1024 pixels
- Calculate the FOV of the camera ...

Photomultiplier Tubes



Trip to Lulin

- Nov 26 Friday, 9:00 am, departure from S4 parking lot
- Bus takes 22-23 people (03-593-2986)
- Lunch at Sui-Li 水里
- ETA 3 pm
- Trail walking ~20-30 min from parking lot to Observatory (0910-267-184)
- No water for shower!
- Bring your own cups
- Bring warm clothes, personal hygienic items, medicines, special food