



Effects of Earth's Atmosphere on Astronomical Observations

1. Limited wavelength ranges (有限的波段)
2. Extinction (消光)
3. Refraction (折射)
4. Curvature (曲率)
5. Atmospheric turbulences (大氣擾動)



1. Limited wavelength ranges

Transparent relatively in optical, radio (and some infrared) → atmospheric “windows”

Optical window: 300 nm to 1.4 μm

O₃ (ozone)
H=25 km


H₂O, CO₂ ...

Radio window : 8 mm to ~15 m

Reflection in
ionosphere
H~100 km

Plasma frequency

Atmospheric Windows




<http://csep10.phys.utk.edu/astr162/lect/light/windows.html>

Q: 波長越短應該吸收越多
但為何 射線吸收少?

大氣吸收的機制:


- 電子躍遷 (electronic transition)
- 分子振動 (vibration; stretching)
- 分子彎擺 (bending)
- 分子轉動 (rotation)

- In addition to photons, the atmosphere also stops energetic charged particles → **cosmic rays** (mainly p^+ , e^-) from space; $E \sim 10^{10}$ - 10^{20} eV



- Atmosphere \approx 1 m of lead
- High-speed particles, if $v \uparrow \uparrow$ → $v > c/n$ (=light speed in the medium, n : refractive index) → ? **erenkov radiation**

Cerenkov radiation



A source emits spherical wavefronts

http://dept.physics.upenn.edu/balloon/cerenkov_radiation.html

- Even beyond earth's atmosphere, there is absorption

→ [dust + gas] in the solar system, in the interstellar medium (星際物質), and circumstellar medium (環星物質)

Dust strong absorption in visual and UV

Gas strong absorption in EUV and soft X-ray



2. Extinction

- **Absorption** (吸收)
photons are "destroyed"
- **Scattering** (散射)
photon energy and direction redistributed
→ Effectively an absorption in the direction to the source

Q: 如何知道某顆星有多亮?

觀測 但，觀測什麼呢？

來自天體的電磁波 → 星際消光

→ 大氣消光 → 望遠鏡收集

→ 偵測器感應 → “讀數”(count)


把這個讀數和「標準星」的讀數相比 → 亮度

例如某顆星的讀數為 1000.0，而利用相同儀器某標準星（視星等為 9.0 等）讀數為 500.0，那麼 ...

Q: 天空為何是藍色的?



Scattering



Size of particles $\sim a$

1. 2 $a \ll (\text{radio}) \lambda$ scattering ?

$I_{\text{scattering}} \propto \lambda^{-4}$ (Rayleigh scattering)

Blue sky

2. 2 $a \gg \lambda$ scattering

Gray sky in a cloudy day!

3. 2 $a \sim \lambda$ (dust, optical) $I_{\text{scattering}} \propto \lambda^{-1}$

⇒ Interstellar reddening (紅化)

3. Refraction

Apparent Altitude	Angle of Refraction
0	35' 21"
1	24' 25"
3	14' 24"
10	05' 18"
30	01' 41"
60	00' 34"
90	00' 00"

- Sun/moon at zenith distance = 90° (refraction $\sim 35'$), but their sizes $\sim 30'$, so when we seen center at horizon, they in fact are below the horizon

4. Curvature


Curvature and refraction can be ignored if zenith distance 45°

If we ignore curvature and refraction,

→ Atmosphere = a series of plane parallel layers, with thickness dz at height z .

$$dF_I = -kF_I \cdot \sec z |dz|$$

where $k(z)$ is the absorption coefficient



忽略大氣折射與曲率，來自天體的光線穿過系列平行大氣層，強度不斷減弱

If F_0 = flux outside atmosphere, then integrating z , we obtain flux at ground level ($z=0$)


$$\Rightarrow F_I(z) = F_0 \exp\left[-\sec z \int_0^z k(z, I) dz\right]$$

At the zenith ($z=0$)

$$F_I(0) = F_0 \exp\left[-\int_0^0 k(z, I) dz\right]$$

$$\Rightarrow \log F_I(z) = \log F_0 + \sec z \cdot \log \frac{F_I(0)}{F_0}$$

Since $m_I = -2.5 \log F_I + \text{const}$



據以修正目標星等

視星等與 $\sec z$ 的關係，並外差到大氣以外的數值

$$m_I = m_0 + \Delta m_0 \cdot \sec z$$

m : observed from ground

m_0 : would have observed outside atmosphere

m_0 : absorption in terms of magnitude at zenith

(cf. $y = b + ax$)

So, m_0 and Δm_0 can be estimated by measuring m (observable) at different $\sec z$ (known).

For 60° , refraction and curvature have to be considered

Define similarly, $m_l = m_0 + \Delta m_0 \cdot M(z)$

where $M(\)$ is the **air mass** = absorption (length) along the curved light-path

For small θ , $M(\theta) \sim \sec \theta$

亦即若天頂角小, $\sec \theta$ 約就是大氣質量

Otherwise determine empirically

⇒ look-up table

But air is not static

→ Try to observe as close to zenith as possible

Apparent zenith distance (degrees)	Air mass
40	01.30
50	01.55
60	02.00
70	02.90
75	03.82
80	05.60
85	10.40
87	15.36
88	19.79

5. Atmospheric Turbulences

大氣擾動 → 有如在游泳池底看世界

Cause the stellar image to

- blink (**scintillation**) variation of air mass along line of sight 一閃一閃亮晶晶
- move around (**'seeing'**) variation of refractive index along line of sight 晃來晃去看不清



“seeing” (視相寧靜度) a point source, after long exposures, is smeared into a ‘seeing’ disk
Typically, seeing (disk) ~ a few arcsec across

www.ee.mtu.edu/~schulz/research/blind_deconvolution.html

Effects of Astronomical Seeing --- M74 (NGC 628)

Distorted image Restored image of HST

http://www.astr.ua.edu/gifimages/m74seeing.html

- Extended sources are not affected as much (averaged out)
e.g. Planets $\sim 10''$ – $30''$ would appear ‘steady’ but stars (point sources) ‘twinkle’.

Usually optical telescopes are seeing-limited

- At radio λ s, with very small-scale turbulences, seeing/scintillation are not important

Radio telescopes are diffraction-limited

i.e., limited by the optics, rather than by atmosphere

Radio λ s are affected by interplanetary and ISM scintillation

To overcome rapid scintillation

⇒ Fast (high-time-resolution) observations

Radio λ s ⇒ Hewish in 1960s tried to study IS scintillation

⇒ Discovered pulsars


Optical λ s ⇒ ‘speckle’ imaging

Speckle Imaging (散斑成像)

連續快速

(0.1 s) 曝光，把大氣

擾動「凍結」




經過影像處理
斑點干涉影像
(左圖)，發現這是顆雙星

一般長期

(40 s) 曝光，大氣擾

動造成星點
影像模糊



Movie

<http://www.ciw.edu/alycia/speckle.html>